

DTIC FILE COPY

2

AD-A197 629

SECURITY CLASSIFICATION OF THIS PAGE

DTIC  
REPORT DOCUMENTATION PAGEForm Approved  
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION U		1b. RESTRICTIVE MARKINGS N/A	
2a SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution Unlimited.	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE N/A		5. MONITORING ORGANIZATION REPORT NUMBER(S) N/A	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Wayne State University		Approved for public release.	
6a NAME OF PERFORMING ORGANIZATION Wayne State University	6b OFFICE SYMBOL (If applicable) N/A	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) Department of Biological Sciences Wayne State University Detroit, Michigan 48202		7b. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0634	
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR04108
		TASK NO. rr1F010	WORK UNIT ACCESSION NO.
11 TITLE (Include Security Classification) (U) Influence of Neuroendocrine Mediators on Phagocyte Function			
12. PERSONAL AUTHOR(S) Howard R. Petty			
13a TYPE OF REPORT Annual	13b TIME COVERED FROM 7/1/87 TO 6/30/88	14. DATE OF REPORT (Year, Month, Day) July 1, 1988	15 PAGE COUNT 14
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
08			
		Stress, Macrophage, Opioid, Adrenergic	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) This research program explores phagocyte responses to neuroendocrine mediators. We have studied the effects of epinephrine, met-enkephalin, forskolin and cAMP analogs on macrophage morphology, spreading and adherence. Cell spreading was quantitated by measuring cell perimeters. Epinephrine decreased macrophage spreading; at $10^{-5}M$ epinephrine the perimeter was $10.4 \pm 0.3 \mu m$ in comparison to $15.0 \pm 1.0 \mu m$ for controls. Epinephrine's action is blocked by propranolol. Met-enkephalin increased macrophage spreading to $18.5 \pm 1.0 \mu m$ at $10^{-8}M$ . Since both catecholamines and opioids are released from chromaffin cells, we have examined the combined effects of both ligands. When macrophages were exposed to $10^{-5}M$ epinephrine and $10^{-8}M$ met-enkephalin, cell morphology and spreading were indistinguishable from that of epinephrine alone. The $\beta$ -adrenergic receptor abrogates the opioid signal(s). These effects may be accounted for by cAMP. Forskolin and cAMP analogs affected cell properties in the same fashion as epinephrine. Dramatic mediator-induced changes in F-actin distribution were noted.			
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION (U)	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. J. A. Maide		22b TELEPHONE (Include Area Code) (202) 696-4055	22c OFFICE SYMBOL ONR


## Influence of Neuroendocrine Mediators on Phagocyte Function

### A. Introduction

It is well known that certain leukocyte receptors evoke synergistic cellular responses. For example, C3b receptor ligation augments immunoglobulin Fc receptor function (18,19). However, phagocyte responses to the ligation of multiple receptors that separately trigger opposing physiologic events have not been thoroughly studied. Two in vitro models of antagonistic combinative interactions are: (1) the histamine and C3bi receptors that promote cell detachment and adherence, respectively (20), and (2) the  $\beta$ -adrenergic and opioid receptors that depress or enhance antibody-dependent phagocytosis (6). It is interesting to note that in both of these combinative systems, the adenylate cyclase-linked receptor abrogates the physiological effects generally ascribed to the second receptor. - In the present study we have examined the individual and combinative effects of epinephrine and met-enkephalin on macrophage morphology, spreading, adherence and microfilaments. Our data: (1) provide evidence indicating that catecholamines down-regulate macrophage activities in the absence or presence of opioids and (2) suggest that elevated cAMP levels play an important role in this activity.

### Effects of Epinephrine, Met-Enkephalin, dbcAMP and Br-cAMP on Cell Morphology

DTIC  
COPY  
INSPECTED  
6

<div style="text-align: center;">  </div>	<div style="text-align: center;"> <p>2025</p> <p>Her</p> </div>
--	---

(Fig. 1b). However,  $10^{-8}$  M met-enkephalin primarily induced the formation of pseudopods and membrane blebs (Fig. 1c). When both of these neuroendocrine mediators are added to cells, their appearance is similar to epinephrine-treated cells (Fig. 1d). Cells treated with dbcAMP or Br-cAMP possessed many surface folds and ruffles, similar to the results obtained with epinephrine (Fig. 1e and f).

The morphology of living macrophages was studied and quantitated by optical microscopy. In comparison to control cells (Fig. 2a), epinephrine at  $10^{-5}$  M (Fig. 2b) did not have an apparent qualitative effect on cell morphology as judged by optical microscopy, although average cell perimeter was diminished (see below). However,  $10^{-8}$  M met-enkephalin caused significant cytoplasmic spreading (see below) and pseudopod formation (Fig. 2c). The combined effect of simultaneous treatment with  $10^{-5}$  M epinephrine and  $10^{-8}$  M met-enkephalin is shown in Figure 2d. The concentrations of these neuroendocrine mediators correspond to the optimal doses found in phagocytosis and spreading assays (see ref. 6). Met-enkephalin is without effect in the presence of epinephrine. The cAMP analogs dbcAMP and Br-cAMP affected cell morphology as judged by bright-field microscopy in a fashion similar to epinephrine (data not shown).

#### Effects of Neuroendocrine Mediators and cAMP Analogues on Cell Spreading and Adherence

To quantitate the effects of neuroendocrine mediators and cAMP analogues on macrophage spreading, the perimeters of macrophages were measured from video images (Fig. 2). Figure 3 shows a dose-response curve illustrating the effect of epinephrine on macrophage spreading. The abscissa is the molar concentration of epinephrine while the ordinate is the percentage of control cell perimeter (the means of all experiments are compared). The measured cell perimeters decrease from  $15.0 \pm 1.0 \mu\text{m}$  for controls to  $10.4 \pm 0.3 \mu\text{m}$  for cells treated with  $10^{-5}$  M epinephrine ( $n = 4$ ,  $P < 0.001$ ). The inhibitory action of epinephrine was blocked by the antagonist propranolol (see below and Fig. 7). The effect of met-enkephalin on macrophage spreading is shown in Figure 4. The measured cell perimeter increases from  $15.0 \pm 1.0 \mu\text{m}$  for controls to  $18.5 \pm 1.0 \mu\text{m}$  for cells treated with  $10^{-8}$  M met-enkephalin ( $n = 5$ ,  $P < 0.001$ ). As qualitatively described by the above micrographs, met-enkephalin induces a significant increase in macrophage spreading whereas epinephrine induces a dramatic decrease in spreading.

Although the above data quantitatively describe the mean cell perimeters during several conditions, they do not illustrate the cell-to-cell variability of the populations. Figure 5 shows the distribution of cell perimeters of macrophages during several experimental conditions. These data represent a compilation of data taken during at least four different trials. Cell perimeters were divided into four groups: 32-43, 46-57, 61-71, and 75-84  $\mu\text{m}$ . These groupings were chosen because they correspond to convenient map reader divisions. The number of cells from all trials within each group were added together then plotted at the ordinate. The effects of epinephrine, met-enkephalin, and the simultaneous addition of both mediators are shown in Figure 5. Although the mean cell perimeter drops 25% in the presence of  $10^{-5}$  M epinephrine, the number of cells in the smallest perimeter group increases over 600%. Similarly,  $10^{-8}$  M met-enkephalin increases mean cell perimeter; the enhancement is most pronounced at the highest perimeter grouping (75-84  $\mu\text{m}$ ). The simultaneous addition of both  $10^{-5}$  M epinephrine and  $10^{-8}$  M met-enkephalin led to a

population distribution indistinguishable from that of epinephrine alone. The combinative effects of epinephrine and met-enkephalin returned to control levels by the addition of propranolol (Fig. 7).

The potential role of cAMP in mediating these changes in spreading behavior was tested using cAMP analogs and forskolin. Figure 6 shows the distributions of cell perimeters for macrophages exposed to 30  $\mu$ M dbcAMP, 30  $\mu$ M Br-cAMP,  $10^{-5}$  M forskolin and matched controls using buffer only. Both cAMP analogs induce a substantial decrease in cell spreading. The effect of dbcAMP could not be reversed by propranolol (Fig. 7). Furthermore, forskolin, a reagent that stimulates endogenous cAMP production, leads to diminished spreading indistinguishable from that of the exogenous analogs. The cell populations of Figures 5 and 6 suggest that: (1) epinephrine screens out the spreading signal of met-enkephalin and (2) this effect may be due to cAMP.

In Figure 7 we show a summary of experiments using the antagonist propranolol. As expected, the addition of  $10^{-5}$  M propranolol has no effect on macrophage spreading ( $99.5 \pm 1.2\%$  of control). In addition, propranolol could not block the ability of dbcAMP to diminish spreading. However, propranolol did inhibit the activity of epinephrine and epinephrine plus met-enkephalin on spreading. Interestingly,  $10^{-5}$  M propranolol diminished the ability of met-enkephalin to augment spreading. The ability of propranolol to influence the activity of opioid receptors was unexpected; participatory factors may include: (1) partial agonist activity of this pharmacologic reagent, (2) a generalized membrane effect, and/or (3) receptor-receptor interactions.

The abilities of epinephrine, met-enkephalin, epinephrine plus met-enkephalin, and dbcAMP to affect RAW264 macrophage adherence were tested. Met-enkephalin at  $10^{-6}$  and  $10^{-8}$  M produced significant and reproducible increases in adherence of  $119 \pm 6.8\%$  and  $112 \pm 5.0\%$  in comparison to controls, respectively ( $n=4$ ;  $P<0.001$  for each). This is consistent with a recent study of neutrophil adherence by Van Epps and Kutvirt (22). Met-enkephalin at other doses and epinephrine and dbcAMP at all doses tested did not influence adherence. No significant change in adherence was found for the combinative experiment using  $10^{-5}$  M epinephrine and  $10^{-8}$  M met-enkephalin ( $n=3$ ;  $92 \pm 16\%$ ).

#### Effects of Neuroendocrine Mediators and cAMP Analogs on Actin Filaments

The ability of epinephrine, met-enkephalin, and cAMP reagents to affect cell morphology and spreading suggests the involvement of cytoskeletal assemblies. To test the effects of these materials on the cytoskeleton, RAW264 macrophages were incubated in the absence or presence of reagents for 30 minutes at  $37^{\circ}\text{C}$ , as described above. Cells were then fixed, extracted, and stained with NBD-phalloidin. Figure 8 shows fluorescence photomicrographs of macrophages after exposure to several conditions. Control and met-enkephalin ( $10^{-8}$  M)-treated cells possessed linear and punctate fluorescent structures (Fig. 8a and c). These structures were particularly apparent in peripheral regions of the cells. In contrast, macrophages treated with epinephrine, epinephrine plus met-enkephalin, dbcAMP, or Br-cAMP demonstrated an intense cortical band of F-actin (Fig. 8b,d,e, and f). This suggests that receptor-cytoskeletal communication, possibly mediated by cAMP, leads to altered cell morphology

and spreading.

#### C. Publications

1. H.R. Petty and K.A. Berg (1988) Combinative Ligand-Receptor Interactions: Epinephrine Depresses RAW264 Macrophage Phagocytosis in the Absence and Presence of Met-Enkephalin. *J. Cell. Physiol.* 134, 281-286.
2. H.R. Petty and S.M. Martin (1988) Combinative Ligand-Receptor Interactions: Effects of cAMP, Epinephrine, and Met-Enkephalin on RAW264 Macrophage Morphology, Spreading, Adherence, and Microfilaments. *J. Cell. Physiol.*, submitted.

#### D. Awards

1. Elected as a Fellow of the American Association for the Advancement of Science
2. Wayne State Fund Research Career Development Chair Award

#### E. Literature Cited

1. Ader, R., ed. (1981) *Psychoneuroimmunology*. Academic Press, N.Y.
2. Blalock, J.E. (1984) *J. Immunol.* 132: 1067-1070.
3. Sanders, V.M., and Munson, A.E. (1985) *Pharmacol. Rev.* 37:229-248.
4. Riley, V. (1981) *Science*. 212:1100-1109.
5. Payan, D.G., Levine, J.D., and Geotzl, E.J. (1984) *J. Immunol.* 132:1601-1604.
6. Petty, H.R., and Berg, K.A. (1988) *J. Cell. Physiol.* 134: 281-286.
7. Koff, W.C., and Dunegan, M.A. (1985) *J. Immunol.* 135:350-354.
8. Heijnen, C.J., Bevers, C., Kavelaars, A., and Ballieux, R.E. (1986) *J. Immunol.* 136:213-216.
9. Gilman, S.C., Schwartz, J.M., Milner, R.J., Bloom, F.E., and Feldman, J.D. (1982) *Proc. Natl. Acad. Sci., USA* 79:4226-4230.
10. Mandler, R.N., Biddison, W.E., Mandler, R., and Sarraf, S.A. (1986) *J. Immunol.* 136:934-939.
11. Van Epps, D.E. and Saland, L. (1984) *J. Immunol.* 132:3046-3053.
12. Abrass, C.K., O'Connor, S.W., Scarpace, P.J., and Abrass, I.B. (1985) *J. Immunol.* 135:1338-1341.
13. Lopker, A., Abood, L.G., Hoss, W., and Lionetti, F.J. (1980) *Biochem. Pharmacol.* 29:1361-1365.
14. Malone, J.D., Richards, M., and Kahn, A.J. (1986) *Proc. Natl. Acad. Sci., USA* 83:3307-3310.
15. Sung, S.-S.J., Young, J. D.-E., Origlio, A.M., Leiple, J.M., Kaback, H.R., and Silverstein, S.C. (1985) *J. Biol. Chem.* 260:13442-13449.
16. Whaley, K., Lappin, D., and Barkas, T. (1981) *Nature* 293:580-583.
17. Bar-Shavit, Z., and Goldman, R. (1986) *Meth. Enzymol.* 132:326-334.
18. Bar-Shavit, Z., Raz, A., and Goldman, R. (1979) *Eur. J. Immunol.* 9:385-391.
19. Ehlenberger, A.G., and Nussenzweig, V. (1977) *J. Exp. Med.* 145:357-371.

20. Francis, J.W., Todd, R.F., Boxer, L.A., and Petty, H.R. (1987) Fed. Proc. 46:1035.
21. Petty, H.R., Hafeman, D.G., and McConnell, H.M. (1981) J. Cell. Biol. 89:223-229.
22. Van Epps, D.E., and Kutvirt, S.L. (1987) J. Neuroimmunol. 15:219-228.

#### FIGURE LEGENDS

Fig. 1. Scanning electron micrographs of macrophages exposed to mediators or cAMP analogs for 30 minutes at 37°C are shown. Cells were prepared for electron microscopy as described in Materials and Methods. Representative micrographs of macrophages treated with: (a) buffer alone (x 2980), (b)  $10^{-5}$  M epinephrine (x 4020), (c)  $10^{-8}$  M met-enkephalin (x 3940), (d)  $10^{-5}$  M epinephrine plus  $10^{-8}$  M met-enkephalin (x 6200), (e) 30  $\mu$ M dbcAMP (x 4020), and (f) 30  $\mu$ M Br-cAMP (x 6200).

Fig. 2. The effects of epinephrine, met-enkephalin, dbcAMP and Br-cAMP on cell morphology are illustrated. Macrophages were allowed to adhere to coverslips for 30 minutes at 37°C in the absence or presence of these substances. Representative bright-field micrographs of macrophages treated with: (a) buffer alone, (b)  $10^{-5}$  M epinephrine, (c)  $10^{-8}$  M met-enkephalin, (d)  $10^{-5}$  M epinephrine and  $10^{-8}$  M met-enkephalin (x 1,200).

Fig. 3. A dose-response curve demonstrating the effect of epinephrine on macrophage spreading is shown. The mean cell perimeter is listed at the ordinate and the molar concentration of epinephrine is given at the abscissa. The mean  $\pm$  s.e.m. of four independent trials are shown.

Fig. 4. A dose-response curve illustrating the effect of met-enkephalin on macrophage spreading is shown. The mean cell perimeter is given at the ordinate; the abscissa lists the molar concentration of met-enkephalin. The mean  $\pm$  s.e.m. of four independent trials are given.

Fig. 5. Analyses of the distribution of macrophage perimeters during four sets of conditions are given. The cell perimeters are distributed between four groupings whose lengths were 32-43, 46-57, 61-71, and 75-84  $\mu$ m. these groups are listed along the abscissa. The total number of cells for all trials is given at the ordinate. The population distributions of cells treated with buffer alone (—),  $10^{-5}$  M epinephrine (---),  $10^{-8}$  M met-enkephalin (...), and  $10^{-5}$  M epinephrine +  $10^{-8}$  M met-enkephalin (\_ \_ \_).

Fig. 6. The distribution of macrophage perimeters in the presence and absence of cAMP analogs is shown. Data are plotted as described in the legend of Fig. 5. The effects of buffer (—), 30  $\mu$ M dbcAMP (...), 30  $\mu$ M Br-cAMP (-----), and  $10^{-5}$  M forskolin (---) are shown (n = 3).

Fig. 7. A bar chart listing the effects of the  $\beta$ -adrenergic antagonist propranolol on macrophage spreading is shown. The percent of the mean control perimeter is shown (mean  $\pm$  s.e.m.). Five sample conditions are given along the abscissa. These conditions are: control,  $10^{-5}$  M epinephrine,  $10^{-8}$  M met-enkephalin,  $10^{-5}$  M epinephrine plus  $10^{-8}$  M met-enkephalin, and 30  $\mu$ M dbcAMP. The open bars represent data in the absence of propranolol; the hatched bars represent identical experiments conducted

in the presence of propranolol (n = 3 to 6; control n = 26 for these trials).

Fig. 8. Fluorescence photomicrographs of RAW264 macrophages labeled with NBD-phalloidin. Cells were exposed to mediators for 30 minutes at 37°C followed by fixation, extraction, and staining. This gallery of photomicrographs includes cells treated with: (a) buffer alone, (b) 10<sup>-5</sup> M epinephrine, (c) 10<sup>-8</sup> M met-enkephalin, (d) 10<sup>-5</sup> M epinephrine plus 10<sup>-8</sup> M met-enkephalin, (e) 30 μM dbcAMP, and (f) 30 μM Br-cAMP. (x 1,000).

Figure 1






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7





Figure 8



## DISTRIBUTION LIST

### Behavioral Immunology Program

Annual, Final and Technical Reports (one copy each except as noted)

#### INVESTIGATORS

Dr. Itamar B. Abrass  
Department of Medicine  
University of Washington  
Harborview Medical Center  
Seattle, WA 98104

Dr. Prince K. Arora  
NIDDK, Bldg. 8, Rm. 111  
National Institutes of Health  
Bethesda, MD 20892

Dr. Andrew S. Baum  
Department of Medical Psychology  
Uniformed Services University  
of Health Sciences, B3050  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

Dr. Charles A. Bowles  
Merrifield Research Lab, Inc.  
P.O. Box 2362  
Merrifield, VA 22116-2362

Dr. Karen Bulloch  
Helicon Foundation  
4622 Sante Fe Street  
San Diego, CA 92109

Dr. Michael D. Cahalan  
Department of Physiology and Biophysics  
University of California, Irving  
Irvine, CA 92717

Dr. Donald A. Chambers  
Health Sciences Center  
University of Illinois at Chicago  
P.O. Box 6998  
Chicago, IL 60680

Dr. Christopher L. Coe  
Department of Psychology  
Harlow Primate Laboratory  
University of Wisconsin  
Madison, IL 53715

Dr. Sheldon Cohen  
Department of Psychology  
Carnegie-Mellon University  
Pittsburgh, PA 15213

Dr. Walla L. Dempsey  
Dept. of Microbiology and  
Immunology  
The Medical College of  
Pennsylvania  
3300 Henry Avenue  
Philadelphia, PA 19129

Dr. David L. Felten  
Department of Anatomy  
University of Rochester  
School of Medicine  
601 Elmwood Avenue  
Rochester, NY 14642

Dr. John F. Hansbrough  
Department of Surgery  
UCSD Medical Center  
225 Dickinson Street  
San Diego, CA 92103

Dr. Robert L. Hunter  
Department of Pathology  
Emory Univ. School of  
Medicine  
WMB 760  
Atlanta, GA 30322

Dr. Terry C. Johnson  
Division of Biology  
Ackert Hall  
Kansas State University  
Manhattan, KS 66506

Dr. Sandra Levy  
University of Pittsburgh  
School of Medicine  
3811 O'Hara Street  
Pittsburgh, PA 15213



Dr. Lester Luborsky  
Department of Psychiatry  
308 Piersol Building/GI  
University of Pennsylvania Hospital  
Philadelphia, PA 19104

Dr. Steven F. Maier  
Department of Psychology  
University of Colorado  
Campus Box 345  
Boulder, CO 80309

Dr. Diana S. Malcolm  
Department of Surgery, USUHS  
Uniformed Services University  
of Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

Dr. Michael H. Melner  
Department of Reproductive Biology  
Oregon Regional Primate Center  
505 N.W. 185th Avenue  
Beaverton, OR 97006

Dr. Vera B. Morhenn  
Department of Dermatology  
Stanford University Medical School  
Stanford, CA 94305

Dr. Jose R. Perez-Polo  
Gail Borden Bldg., Rm. 436  
University of Texas Medical Branch  
Galveston, TX 77550-2777

Dr. Howard R. Petty  
Department of Biological Sciences  
Wayne State University  
Detroit, MI 48202

Dr. Bruce S. Rabin  
Clinical Immunopathology  
Childrens Hospital  
University of Pittsburgh School of  
Medicine  
Pittsburgh, PA 15213

Dr. Seymour Reichlin  
Director, Clinical Study Unit  
New England Medical Center Hospitals, Inc.  
171 Harrison Avenue  
Boston, MA 02111

Dr. Eric M. Smith  
Department of Psychiatry  
University of Texas  
Medical Branch  
Galveston, TX 77550

Dr. Ross R. Vickers, Jr.  
Naval Health Research Ctr.  
Bldg. 346  
P.O. Box 85122  
San Diego, CA 92138

Annual, Final and Technical Reports (one copy each except as noted)

ADMINISTRATORS

Dr. Jeannine A. Majde, Code 1141SB (2 copies)  
Scientific Officer, Immunology Program  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000

Program Manager  
Biological/Human Factors Division  
Office of Naval Research, Code 125  
800 N. Quincy Street  
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

Program Manager  
Support Technology Directorate  
Office of Naval Technology, Code 223  
800 N. Quincy Street  
Arlington, VA 22217-5000

Administrative Contracting Officer  
ONR Resident Representative  
(address varies - obtain from business office)

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

Commanding Officer  
Naval Medical Center  
Washington, DC 20372

Commander  
USAMRIID  
Fort Detrick  
Frederick, MD 21701

Commanding Officer  
Naval Medical Research & Development Command  
National Naval Medical Center  
Bethesda, MD 20814

Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332

Director, Infectious Diseases Program Center  
Naval Medical Research Institute  
National Naval Medical Center  
Bethesda, MD 20814

Library  
Armed Forces Radiation Research  
Institute  
Bethesda, MD 20814-5145

Commander  
Chemical and Biological Sciences Division  
Army Research Office, P.O. Box 12211  
Research Triangle Park, NC 27709

Commander  
U.S. Army Research and Development Command  
Attn: SGRD-PLA  
Fort Detrick  
Frederick, MD 21701

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)  
Attn: Technical Information Division, Code 2627  
Washington, DC 20375